

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/003061

International filing date: 22 March 2005 (22.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: GB
Number: 0415043.9
Filing date: 05 July 2004 (05.07.2004)

Date of receipt at the International Bureau: 23 May 2005 (23.05.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



PCT/EP 200 5 / 0 0 3 0 6 1
5 / 0 0 3 0 6 1

03. 05. 2005



INVESTOR IN PEOPLE


The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 14 April 2005

Patents Form 1/77

Patents Act 1977
(Rule 16)05JUL04 E90817-1 010086
P01/7700 0.00-0415043.9 ACCOUNT CHA

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

THE PATENT OFFICE
RM
- 5 JUL 2004
RECEIVED BY FAX

1. Your reference

CBG 00230/GB/P1

2. Patent application number

(The Patent Office will fill in this part)

0415043.9

3. Full name, address and postcode of the or of each applicant: (underline all surnames)

Complex Biosystems GmbH
Im Neuenheimer Feld 584
D-69120 Heidelberg
Germany

Patents ADP number (if you know it)

8903817001

If the applicant is a corporate body, give the country/state of its incorporation

Germany

4. Title of the invention

Prodrug Linker

5. Name of your agent (if you have one)

Martin Hyden

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Rouse Patents
Windsor House
Cornwall Road
Harrogate
HG1 2FW
United Kingdom

Patents ADP number (if you know it)

8661159001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

Yes

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 65

Claim(s)

Abstract

Drawing(s) 6 only RM

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents
(please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

MARTIN HYDEN

05/07/04

12. Name and daytime telephone number of person to contact in the United Kingdom

Jane Polizzi/Anna Whitehead 01423 850800

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Prodrug Linker

Field

- 5 The present invention is directed to polymeric prodrugs having reversible linkages to amino groups of biologically active entities such as peptides, proteins, natural products or synthetic chemical compounds.

Background

10

Polymers used in drug delivery systems are of interest because the therapeutic efficacy of both many low molecular weight and macromolecular medicinal agents may be significantly improved by combining them with polymers.

- 15 For example, covalent modification of biological molecules with poly(ethylene glycol) (PEG) has been extensively studied since the late 1970s. So-called PEGylated proteins have shown improved therapeutic efficacy by increasing solubility, reducing immunogenicity, and increasing circulation half-life in vivo due to reduced renal clearance and proteolysis by enzymes (Caliceti P., Veronese F.M., Adv. Drug Deliv. Rev. 2003, 55, 1261-1277).
- 20

- Many small molecule medicinal agents, like alkaloids and anti-tumor agents, show low solubility in aqueous fluids. One way to solubilize these small molecule compounds is to conjugate them to hydrophilic polymers. A variety of water-soluble polymers, such as human serum albumin, dextran, lectins, poly(ethylene glycol), poly(styrene-co-maleic anhydride), poly(N-hydroxypropylmethacrylamide), poly(divinyl ether-co-maleic anhydride), hyaluronic acid have been described for this purpose.
- 25

- 30 Depot formulations generating long-acting release profiles have been described using non-covalent drug encapsulation. Typically, a drug is mixed with polymer material and processed in such fashion, that the drug is captured in the polymer interior.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Resulting polymer-protein aggregates may be shaped as microparticles or gels. Drug release occurs if the polymer swells or degradation of the polymer allows for diffusion of the drug to the exterior. Degradation processes may be autohydrolytic or enzyme-catalyzed. Marketed drugs based on non-covalent polymer encapsulation are Lupron Depot and Nutropin Depot. A disadvantage of this approach is that in order to prevent uncontrolled, burst-type release of the non-covalently bound drug, encapsulation has to be highly efficient, requiring strong van der Waals contacts, frequently mediated through hydrophobic moieties, and many conformationally sensitive therapeutics such as proteins or peptides are rendered dysfunctional during the encapsulation process and/or during subsequent storage. Furthermore, dependence of the release mechanism upon biodegradation may cause interpatient variability.

A major challenge in cancer therapy is to selectively target cytotoxic agents to tumor cells. A promising method to accumulate small molecule anticancer agents in tumor tissue and decrease undesirable side effects of these agents is the attachment of the cytotoxin to a macromolecular carrier. The passive targeting of polymeric drug conjugates to tumors is based on the so-called enhanced permeability and retention effect (EPR) as described by Matsumura, Y. and Maeda, H., in *Cancer Res.*, 1986, vol 6, pp 6387-6392. As a result, several polymer-drug conjugates have entered clinical trial as anticancer agents.

However, many medicinal agents are inactive or show decreased biological activity when a polymer is covalently conjugated to the drug molecule. In these cases, a prodrug approach may be applied in such a fashion that the medicinal agent is released from the polymeric carrier in vivo to regain its biological activity. Prodrugs may be defined as therapeutic agents that are inactive per se but are predictably transformed into active metabolites (see B. Testa, J.M. Mayer in *Hydrolysis in Drug and Prodrug Metabolism*, Wiley-VCH, 2003, page 4). The reduced biological activity of the prodrug as compared to the released drug is of advantage if a slow or controlled release of the drug is desired. In this case, a relatively large amount of prodrug may be administered without concomitant side effects and the risk of overdosing. Release of the drug occurs over time, thereby reducing the necessity of repeated and frequent administration of the drug.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Definitions established by IUPAC

(as given under <http://www.chem.qmul.ac.uk/iupac/medchem/> (accessed on 8 March 2004))

5 Prodrug

A prodrug is any compound that undergoes biotransformation before exhibiting its pharmacological effects. Prodrugs can thus be viewed as drugs containing specialized non-toxic protective groups used in a transient manner to alter or to eliminate undesirable properties in the parent molecule.

10

Double prodrug (or pro-prodrug)

A double prodrug is a biologically inactive molecule which is transformed *in vivo* in two steps (enzymatically and/or chemically) to the active species.

15

Carrier-linked prodrug (Carrier prodrug)

A carrier-linked prodrug is a prodrug that contains a temporary linkage of a given active substance with a transient carrier group that produces improved physicochemical or pharmacokinetic properties and that can be easily removed *in vivo*, usually by a hydrolytic cleavage. This is shown graphically in Fig. 1.

20

Cascade prodrug

A cascade prodrug is a prodrug for which the cleavage of the carrier group becomes effective only after unmasking an activating group. This is shown graphically in Fig.

25 2.

Bioprecursor prodrug

A bioprecursor prodrug is a prodrug that does not imply the linkage to a carrier group, but results from a molecular modification of the active principle itself. This modification generates a new compound, able to be transformed metabolically or chemically, the resulting compound being the active principle.

30

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Biotransformation

Biotransformation is the chemical conversion of substances by living organisms or enzyme preparations.

- 5 This invention applies in particular to carrier prodrugs and cascade prodrugs. In polymeric prodrugs, the drugs are often linked to a polymeric carrier moiety by a labile bridge formed between the carrier moiety and a hydroxy, amino or carboxy group of the drug molecule.
- 10 Cleavage of a carrier prodrug generates a molecular entity (drug) of increased bioactivity and at least one side product, the carrier. This side product may be biologically inert (for instance PEG) or may have targeting properties (for instance antibodies). After cleavage, the bioactive entity will reveal at least one previously conjugated and thereby masked or protected functional group, and the presence of this
- 15 group typically contributes to the bioactivity. Numerous macromolecular prodrugs are described in the literature where the macromolecular carrier is linked via a labile ester group to the medicinal agent (e.g. Y. Luo, MR Ziebell, GD Prestwich, "A Hyaluronic Acid - Taxol Antitumor Bioconjugate Targeted to Cancer Cells", Biomacromolecules 2000, 1, 208-218, J Cheng et al, Synthesis of Linear, beta-Cyclodextrin Based
- 20 Polymers and Their Camptothecin Conjugates, Bioconjugate Chem. 2003, 14, 1007-1017, R. Bhatt et al, Synthesis and in Vivo Antitumor Activity of Poly(L-glutamic acid) Conjugates of 20(S)-Camptothecin, J. Med. Chem. 2003, 46, 190-193 ; R.B. Greenwald, A. Pendri, C.D. Conover, H. Zhao, Y.H. Choe, A. Martinez, K. Shum, S. Guan, J. Med. Chem., 1999, 42, 3657-3667; B. Testa, J.M. Mayer in Hydrolysis in
- 25 Drug and Prodrug Metabolism, Wiley-VCH, 2003, Chapter 8) In these cases, the conjugated functional group of the bioactive entity is a hydroxyl group or a carboxylic acid.
- 30 Especially for biomacromolecules but also for small molecule polymer prodrugs, it may be desirable to link the macromolecular carrier to amino groups (i.e. N-terminus or lysine amino groups of proteins) of the bioactive entity, as is shown in Fig. 3. This will be the case if masking the drug's bioactivity requires conjugation of a certain amino group of the bioactive entity, for instance an amino group located in an active center or a region or epitope involved in receptor binding. Also, during preparation of

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

the prodrug, amino groups may be more chemoselectively addressed and serve as a better handle for conjugating carrier and drug because of their greater nucleophilicity as compared to hydroxylic or phenolic groups. This is particularly true for proteins which may contain a great variety of different reactive functionalities. In this case
5 non-selective conjugation reactions lead to undesired product mixtures which require extensive characterization or purification and may decrease reaction yield and therapeutic efficiency of the product.

10 Prodrug activation may occur by enzymatic or non-enzymatic cleavage of the labile bridge between the carrier and the drug molecule, or a sequential combination of both, i.e. an enzymatic step followed by a nonenzymatic rearrangement.

In an enzyme-free in vitro environment such as an aqueous buffer solution, almost
15 any prodrug will undergo hydrolysis if labile linkages such as ester or amide bonds are present in the molecule. This rate of hydrolysis may be very slow and not therapeutically useful. In an in vivo environment, esterases or amidases are typically present and may cause significant catalytic acceleration of the kinetics of hydrolysis from twofold up to several orders of magnitude (R.B. Greenwald et al. J.Med.Chem.
20 1999, 42 (18), 3857-3867).

The labile bond may be engineered for enzyme selectivity for organ or cell-type targeting. Targeted release of the bioactive entity is effected, if an enzyme, that selectively cleaves the linkage, is specifically present in the organ or cell type chosen
25 for treatment. Several examples were published for the prodrug activation of amine containing drugs by specific enzymes. In these cases, targeted release was the objective. G. Cavallaro et al., Bioconjugate Chem. 2001, 12, 143-151 described the enzymatic release of an antitumoral agent by the protease plasmin. Cytarabin was coupled via the tripeptide sequence D-Val-Leu-Lys to the polymer alpha, beta-
30 poly(N-hydroxyethyl)-DL-aspartamide (PHEA). Enzymatic release of cytarabin was effected by the protease plasmin which concentration is relatively high in various kinds of tumor mass.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

- Further examples for antitumoral polymeric prodrugs activated by specific enzymes like beta lactamase (R. Satchi-Fainaro et al., *Bioconjugate Chem.* 2003, 14, 797-804) and cysteine proteases like cathepsin B (R. Duncan et al. *J. Contr. Release* 2001, 74, 135-146) were described. Wiwattanapatapee et al. (2003) described a dendrimer
- 5 prodrug for colonic delivery of 5-aminosalicylic acid. The drug molecule was conjugated by an azo bond to generation 3 PAMAM dendrimer. 5-aminosalicylic acid is released in the colon by a bacterial enzyme called azo reductase (W. R. Wiwattanapatapee, L. Lomlim, K. Saramunee, *J. Controlled Release*, 2003, 88: 1-9).
- 10 A.J. Garman et al. (A.J. Garman, S.B. Kalindjian, *FEBS Lett.* 1987, 223 (2), 361-365 1987) used PEG5000-maleic anhydride for the reversible modification of amino groups in tissue-type plasminogen activator and urokinase. Regeneration of functional enzyme from PEG-uPA conjugate upon incubation at pH 7.4 buffer by cleavage of the maleamic acid linkage followed first order kinetics with a half-life of 6.1 h. The
- 15 prodrug cleavage was not investigated in the presence of enzymes, and it can be expected – as explained above – that in vivo proteases will significantly contribute to the cleavage of the labile amide linkage. A further disadvantage of this linkage is the lack of stability of the conjugate at lower pH values that limits this approach to active agents stable at basic pH values as purification of the active agent polymer conjugate
- 20 has to be performed under basic conditions.
- R.B. Greenwald, A. Pendri, C.D. Conover, H. Zhao, Y.H. Choe, A. Martinez, K. Shum, S. Guan, *J. Med. Chem.*, 1999, 42, 3657-3667 & PCT Patent Application WO-A-99/30727 described a methodology for synthesizing poly(ethylene glycol) prodrugs
- 25 of amino-containing small molecule compounds based on 1,4- or 1,6-benzyl elimination. In this approach the amino group of the drug molecule is linked via a carbamate group to a PEGylated benzyl moiety. The poly(ethylene glycol) was attached to the benzyl group by ester, carbonate, carbamate, or amide bonds. The release of PEG from the drug molecule relied on the enzymatic cleavage of these
- 30 masking groups. The cleavage of the release-triggering masking group is followed in this approach by the classical and rapid 1,4- or 1,6-benzyl elimination. As a consequence the highly reactive and potentially toxic quinone methide or quinonimine methide is liberated.

Ref: CB00230/GB/F1
Applicant: Complex Biosystems GmbH

5 July 2004

This is an example for a cascade carrier prodrug where the masking group is identical with the carrier and is shown diagrammatically in Fig. 4.

5 This linker system was also used for releasable poly(ethylene glycol) conjugates of proteins (S. Lee, R.B. Greenwald et al. Bioconj. Chem. 2001, 12 (2), 163-169).
Lysozyme was used as model protein because it loses its activity when PEGylation takes place on the epsilon-amino group of lysine residues. Various amounts of PEG linker were conjugated to the protein. Regeneration of native protein from the PEG conjugates occurred by enzymatic cleavage in rat plasma or in non-physiological high
10 pH buffer.

Greenwald et al. published in 2000 a poly(ethylene glycol) drug delivery system of amino-containing prodrugs based on trimethyl lock lactonization (R.B. Greenwald et al. J.Med.Chem. 2000, 43(3), 457-487; PCT Patent Application No. WO-A-
15 02/089789). In this prodrug system substituted o-hydroxyphenyl-dimethylpropionic acid is coupled to amino groups of drug molecules by an amide bond. The hydroxy group is linked to PEG by an ester, carbonate, or carbamate group. The rate determining step in drug release is the enzymatic cleavage of these functional groups followed by fast amide cleavage by lactonization, liberating a potentially toxic
20 aromatic lactone side product.

A common disadvantage of the linker systems described by Greenwald is the release of a potentially toxic aromatic small molecule side product after cleavage of these prodrug linker molecules.

25 More recently, R.B. Greenwald et al. (Greenwald et al. J. Med.Chem. 2004, 47, 726-734) described a PEG prodrug system based on bis-(N-2-hydroxyethyl)glycin amide (bicin amide) linker. In this system two PEG molecules are linked to a bicin molecule coupled to an amino group of the drug molecule. The first two steps in prodrug
30 activation is the enzymatic cleavage of both PEG molecules. Different linkages between PEG and bicin are described resulting in different prodrug activation kinetics. The main disadvantage of this system is the slow hydrolysis rate of bicin amide conjugated to the drug molecule ($t_{1/2} = 3$ h in phosphate buffer) resulting in the release of a bicin-modified prodrug intermediate that may show different

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

pharmacokinetic and pharmacodynamic properties compared to the parent drug molecule.

5 The in vivo release of stable linker-drug intermediates or potentially toxic side products can be avoided, if the carrier is not serving as a masking group and the activating group is coupled to the carrier by means of a stable bond. Antczak et al. (Bioorg Med Chem 9 (2001) 2843-48) describe a reagent which forms the basis for a cascade prodrug system for amine-containing drug molecules. In this approach an antibody serves as carrier, a stable bond connects the antibody to an activating moiety, 10 carrying an enzymatically cleavable masking group. Upon enzymatic removal of the masking group, a second labile bond cleaves and releases the drug compound, as shown in Fig. 5.

15 A drawback of predominantly enzymatic cleavage is interpatient variability. Enzyme levels may differ significantly between individuals resulting in biological variation of prodrug activation by enzymatic cleavage. Enzyme levels may also vary depending on the site of administration, for instance it is known that in the case of subcutaneous injection, certain areas of the body yield more predictable therapeutic effects than others. To reduce this unpredictable effect, non-enzymatic cleavage or intramolecular 20 catalysis is of particular interest (see, for example, B. Testa, J.M. Mayer in Hydrolysis in Drug and Prodrug Metabolism, Wiley-VCH, 2003, page 5).

Using enzyme specificity to control both targeting and release kinetics is a compromise. Enzyme-dependent release results in interpatient variability. Targeting 25 through enzyme selectivity is of lower efficiency than for instance targeting through antibodies and may be achieved by more reliable means for instance by introducing certain structural features into the polymer moiety of the prodrug.



30 For these reasons, there is a need to provide novel linker and/or carrier technologies for forming polymeric prodrugs of amine containing active agents in order to overcome the limitations of the described polymeric prodrugs.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


Detailed description of the invention

The present invention addresses the disadvantages described above. The invention provides for a polymeric prodrug of Formula Ia or Ib.



wherein Y_1 to Y_5 , R_1 to R_4 , T , X , W , Nu and Ar are defined below:

Native drug release is effected by a two step mechanism. The first step is the rate-determining cleavage of the masking group:




from the polymeric prodrug in vivo.

30 As described above, cleavage of the masking moiety may be mediated by an enzymatic or a non-enzymatic step, such as pH-dependent hydrolysis or intramolecular cyclization. In the preferred embodiment of the invention, cleavage is effected non-enzymatically by intramolecular cyclization or catalysis. The half-life of the cleavage kinetics in aqueous buffer of pH 7.4 at 37 °C of the masking group

Ref: CBG00230/GB/FI
Applicant: Complex Biosystems GmbH


5 July 2004

according to the present invention is preferably between 1 h to 6 month, more preferably between 1 d to 3 months, and most preferably between 1 d to 2 months. The second and final step in release of the regenerated native drug is the fast, spontaneous and irreversible so-called 1,4- or 1,6 or 1,(4+2n) with $n = 2, 3, 4$ or higher elimination of the



moiety of the remaining polymeric prodrug of formula Ia or formula Ib, respectively.

This mechanism of native drug release from a polymeric prodrug triggered by hydrolytic cleavage of the masking group followed by a 1,6-elimination step of the activating group is exemplified by an example of a polymeric prodrug according to the present invention.



Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Definition of Y₁ to Y₅, R1 to R4, T, X, W, Nu and Ar in formula Ia or Ib

T is D or A

5 and D is a residue of an amine-containing biologically active material including but not limited to small molecule bioactive agents or biopolymers like proteins, polypeptides and oligonucleotides (RNA, DNA), peptide nucleic acids (PNA),

and A is a leaving group.

10

Non-limiting examples of suitable leaving groups A include chloride, bromide, fluoride, nitrophenoxy, imidazolyl, N-hydroxysuccinimidyl, N-hydroxybenzotriazolyl, N-hydroxyazobenzotriazolyl, pentafluorophenoxy, N-hydroxysulfosuccinimidyl, or any other leaving group known by those skilled in the art.

15

Suitable organic small molecule bioactive drugs include, without limitation, moieties such as central nervous system-active agents, anti-infective, anti-neoplastic, antibacterial, anti-fungal, analgesic, contraceptive, anti-inflammatory, steroidal, vasodilating, vasoconstricting, and cardiovascular agents with at least one primary or secondary amino group. Non-exclusive examples of such compounds are daunorubicin, doxorubicin, idarubicin, mitoxantron, aminoglutethimide, amantadine, diaphenylsulfon, ethambutol, sulfadiazin, sulfamerazin, sulfamethoxazol, sulfalen, cinafloxacin, moxifloxacin, ciprofloxacin, enoxacin, norfloxacin, neomycin B, sprectinomycin, kanamycin A, meropenem, dopamin, dobutamin, lisinopril, serotonin, carbutamid, acivicin, etc.

20

25

Suitable proteins and polypeptides having at least one free amino group include but are not limited to ACTH, adenosine deaminase, agalsidase, albumin, alfa-I antitrypsin (AAT), alfa-I proteinase inhibitor (API), alteplase, anistreplase, anicod serine protease, antibodies (monoclonal or polyclonal, and fragments or fusions), antithrombin III, antitrypsins, aprotinin, asparaginases, biphalin, bone-morphogenic proteins, calcitonin (salmon), collagenase, DNase, endorphins, enfuvirtide, enkephalins, erythropoietins, factor VIIa, factor VIII, factor VIIla, factor IX,

30

Ref: CBO00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

fibrinolysin, fusion proteins, follicle-stimulating hormones, granulocyte colony stimulating factor (G-CSF), galactosidase, glucagon, glucocerebrosidase, granulocyte macrophage colony stimulating factor (GM-CSF), phospholipase-activating protein (PLAP), gonadotropin chorionic (hCG), hemoglobins, hepatitis B vaccines, hirudin,
5 hyaluronidases, iduronidase, immune globulins, influenza vaccines, interleukins (1 alfa, 1 beta, 2, 3, 4, 6, 10, 11, 12), IL-1 receptor antagonist (rhIL-1ra), insulins, interferons (alfa 2a, alfa 2b, alfa 2c, beta 1a, beta 1b, gamma 1a, gamma 1b), keratinocyte growth factor (KGF), transforming growth factors, lactase, leuprolide, levothyroxine, luteinizing hormone, lyme vaccine, natriuretic peptide, pancrelipase,
10 papain, parathyroid hormone, PDGF, pepsin, platelet activating factor acetylhydrolase (PAF-AH), prolactin, protein C, octreotide, secretin, somatostatin, superoxide dismutase (SOD), somatropins (growth hormone), somatostatin, streptokinase, sucrase, tetanus toxin fragment, tilactase, thrombins, thymosin, thyroid stimulating hormone, thyrotropin, tumor necrosis factor (TNF), TNF receptor-IgG Fc, tissue plasminogen
15 activator (tPA), TSH, urate oxidase, urokinase, vaccines, plant proteins such as lectins and ricins.

Also included herein is any synthetic polypeptide or any portion of a polypeptide with in vivo bioactivity. Furthermore, proteins prepared by recombinant DNA
20 methodologies including mutant versions of aforementioned proteins, antibody fragments, single chain binding proteins, catalytic antibodies and fusion proteins are included.

Preferred proteins are antibodies, calcitonin, G-CSF, GM-CSF, erythropoietins,
25 hemoglobins, interleukins, insulins, interferons, SOD, somatropin, TNF, TNF-receptor-IgG Fc.

X is a spacer moiety such as R5-Y6

30 Y₁, Y₂ can each be either O, S, or NR6, independently of each other.

Y₃, Y₅ can each be either O or S, independently of each other.

Y₄ is O, NR6, or -C(R7)(R8)-

Y₆ is O, S, NR6, succinimide, maleimide, unsaturated carbon-carbon bonds or any heteratom containing a free electron pair or is not present.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

R2 and R3 are selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryls, cyano, nitro, halogen, carboxy, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, etc.

5

The term "heteroalkyl" in the context of the present invention denotes (linear, cyclical or branched) alkyl chains where the alkyl chains contain one or more heteroatoms, selected independently from O, S, N, P, Si, etc. or groups, selected independently from amide, ester, phosphonate ester, phosphate ester, double or triple bonds, carbamate, urea, thiourea, thiocarbamate, oxime, etc.

10

Each R4 substitution on Ar may be the same or different and is selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryl, substituted aryl, substituted or non-substituted heteroaryl, substituted or non-substituted linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy, heteroaryloxy, etc.

15

R4 is selected preferably from small substituents such as hydrogen, methyl, ethyl, ethoxy, methoxy, and other C1 to C6 linear, cyclical or branched alkyls and heteroalkyls.

20

n is zero or a positive integer.

R7 and R8 are selected independently from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryls, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, cyano, halogen, etc.

25

R5 is selected from substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryls, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, etc.

30

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

R6 is selected from hydrogen, substituted or non-substituted linear, branched or cyclical alkyl or heteroalkyl, aryls, substituted aryls, substituted or non-substituted heteroaryls, etc.

5 R1 is a polymer.

Non-limiting examples for suitable polymers are polyalkyloxy-based polymers like poly(propylene glycol) or poly(ethylene glycol), dextran, chitosan, hyaluronic acid and derivatives, alginate, xylan, mannan, carrageenan, agarose, cellulose, starch, hydroxyethyl starch (HES) and other carbohydrate-based polymers, poly(vinyl
10 alcohols), poly(oxazolines), poly(anhydrides), poly(ortho esters), poly(carbonates), poly(urethanes), poly(acrylic acids), poly(acrylamides) such as poly(hydroxypropylmethacrylamide) (HMPA), poly(acrylates), poly(methacrylates) like poly(hydroxyethylmethacrylate), poly(organophosphazenes), poly(siloxanes), poly(vinylpyrrolidone), poly(cyanoacrylates), poly(esters) such as poly(lactic acid) or
15 poly(glycolic acids), poly(iminocarbonates), poly(amino acids) such as poly(glutamic acid), collagen, gelatin, copolymers, grafted copolymers, cross-linked polymers, hydrogels, and block copolymers from the above listed polymers.

Hydrogels may be defined as three-dimensional, hydrophilic or amphiphilic polymeric networks imbibing large quantities of water. The networks are composed of
20 homopolymers or copolymers, are insoluble due to the presence of covalent chemical or physical (ionic, hydrophobic interactions, entanglements) crosslinks. The crosslinks provide the network structure and physical integrity. Hydrogels exhibit a thermodynamic compatibility with water which allows them to swell in aqueous media. (Lit.: N.A. Peppas, P. Bures, W. Leobandung, H. Ichikawa, Hydrogels in
25 pharmaceutical formulations, Eur. J. Pharm. Biopharm. 2000, 50, 27-46). The chains of the network are connected in such a fashion that pores exist and that a substantial fraction of these pores are of dimensions 1...1000 nm. By selecting certain polymerization conditions, the hydrogel may be obtained in the form of an amorphous gel or as beaded resin. Such soft beads may have a diameter of 1...1000 micrometer.
30 Hydrogels may be synthesized from the polymers and copolymers listed above and physically cross-linked or chemically crosslinked by radical, anionic or cationic polymerization, by chemical reactions like condensation or addition reactions as described in W.E. Hennink and C.F. van Nostrum, Adv. Drug Del. Rev. 2002, 54, 13-36.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Further examples include branched and hyperbranched polymers. Examples for such polymers include dendrimers and other dense star polymers. (R. Esfand, D.A. Tomalia, Drug Discov Today, 2001, 6(8), 427-436; P.M. Heegaard, U. Boas, Chem. Soc. Rev. 2004 (33(1), 43-63; S.M. Grayson, J.M. Frechet, Chem. Rev. 2001, 101 (12), 3819-3868).

R1 can also be a biopolymer like a protein. Non-limiting examples include albumin, antibodies, fibrin, casein, and other plasma proteins.

Each polymer can carry one or more biologically active substances linked to the polymer by the prodrug linker described herein or any other linker known to the person skilled in the art. The polymers may have further substituents and may be functionalized for attachment to the spacer moiety X. Non-limiting examples of such functional groups comprise carboxylic acid and activated derivatives, amino, maleimide, thiol, sulfonic acid and derivatives, carbonate and derivatives, carbamate and derivatives, hydroxyl, aldehyde, ketone, hydrazine, isocyanate, isothiocyanate, phosphoric acid and derivatives, phosphonic acid and derivatives, haloacetyl, alkyl halides, acryloyl, arylating agents like aryl fluorides, hydroxylamine, disulfides like pyridyl disulfide, vinyl sulfone, vinyl ketone, diazoalkanes, diazoacetyl compounds, epoxide, oxirane, and aziridine.

Preferred functional groups for the polymer include but are not limited to thiol, maleimide, amino, carboxylic acid and derivatives, carbonate and derivatives, carbamate and derivatives, aldehyde, and haloacetyl.

Especially preferred functional groups include thiol, maleimide, amino, carboxylic acid and derivatives, carbamate and derivatives, and carbonate and derivatives thereof.

Non-limiting examples for suitable bonds or groups formed between X and R1 include disulfide, S-succinimido, amide, amino, carboxylic ester, sulfonamide, carbamate, carbonate, ether, oxime, hydrazone, urea, thiourca, phosphate, phosphonate, etc.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


Preferred bonds or groups formed between X and R1 comprise S-succinimido, amide, carbamate, and urea.

- 5 Preferably, the polymers are well hydrated, degradable or excretable, nontoxic and non-immunogenic in mammals. Preferred polymers include polyalkoxy-based polymers like polyethylene glycol and polyethylene glycol reagents as those described in Nektar Inc. 2003 catalog "Nektar Molecule Engineering – Polyethylene Glycol and Derivatives for Advanced PEGylation" and branched, hyperbranched,
10 cross-linked polymers and hydrogels.

- W is selected from substituted or non-substituted linear, branched or cyclical alkyl, aryls, substituted aryls, substituted or non-substituted linear, branched or cyclical heteroalkyl, substituted or nonsubstituted heteroaryls, substituted or non-substituted
15 linear, branched, or cyclical alkoxy, substituted or non-substituted linear, branched, or cyclical heteroalkyloxy, aryloxy, heteroaryloxy, cyano, carboxy, carboxyalkyl, alkylcarbonyl, carboxamidoalkyl, etc.

W is selected preferably from non-toxic substituted or non-substituted linear, branched or cyclical alkyls or heteroalkyls.

20



At least one Nu is present in Nu-W.


- 25 Nu is a nucleophile that can perform a nucleophilic attack at the carbonyl carbon of $Y_4 \text{---} \overset{Y_1}{\parallel} \text{---} Y_2$ and thus catalyse the cleavage of the masking group by intramolecular catalysis or cyclization. Nucleophiles are any functional groups or heteroatoms carrying at least one free electron pair. Preferred nucleophiles include primary,
secondary and tertiary amino groups, thiol, carboxylic acid, hydroxylamine,
30 hydrazine. In order to effectively catalyse the cleavage of the masking group, the spacing between the nucleophile Nu and Y_2 is preferably between three and fifteen atoms. More preferably, the spacing between Nu and Y_2 is between four and eight atoms. The at least one nucleophile Nu may be attached anywhere to W (e.g. at the terminus or in the middle of W) or may be part of W.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

An especially preferred variation of $Y_4 \overset{Y_1}{\parallel} Y_2$ is $NR6-\overset{O}{\parallel}-O$ and R6 may also be Nu-W and is not hydrogen. In this case, preferred variations of each Nu-W are selected independently from substructures according to the following substructure:

5



R9, R10, R11, and R12 are selected independently from hydrogen, substituted or non-substituted alkyl or heteroalkyl, substituted or non-substituted aryl or heteroaryl.

10

Preferably, R9, R10, R11, and R12 are selected independently from hydrogen, substituted or non-substituted alkyl.

15

Figure 8 shows an example according to formula Ia or Ib where cleavage of the masking group is by intramolecular cyclization.

In cases where Nu only catalyses the cleavage of the masking group by intramolecular catalysis, no cyclical product of the masking group is formed.

20

Surprisingly it was found, that the masking group can modify irreversibly the amine containing biologically active moiety when the nucleophile Nu is absent in the masking group. As shown in the example section, during release of the bioactive moiety insulin from a polymeric prodrug with a pentanoyl masking group which is not part of the present invention, approximately 30 % of the insulin molecule was modified with the masking group by acyl transfer. The mechanism of an example of this modification where D contains an additional free amino group that serve as nucleophile for acyl transfer is shown in figure 9.


25

Ar of formula Ia or Ib is a multi-substituted aromatic hydrocarbon or a multi-substituted aromatic heterocycle. To be aromatic, the number of pi electrons must satisfy the Hückel rule ($4n+2$) and the cycle has to be planar. A huge variety of compounds satisfy these criteria and thus are suitable as Ar in formula Ia or Ib. Non-limiting preferred aromatic moieties include:

30

Ref: CBG00230/CB/P1
Applicant: Complex Biosystems GmbH

5 July 2004



20 wherein W is O, N, or S, independent from each other.

25 Y_2 and $\begin{array}{c} R1 \\ | \\ X \\ | \\ R3 \end{array} \begin{array}{c} Y5 \\ || \\ T \end{array}$ in formula Ia or $\begin{array}{c} R2 \\ | \\ Y3 \end{array} \begin{array}{c} Y5 \\ || \\ T \end{array}$ in formula Ib have to be arranged on the aromatic ring in such a fashion that a 1,4- or 1,6- or 1,(4+2n), with n= 2, 3, 4 and higher, elimination can take place (see above). For example, in the case of a 6-membered ring, the substituents have to be arranged *ortho* or *para*.

30 Preferred moieties for Ar are mono- and bicyclic aromatic hydrocarbons or aromatic heterocycles.

Especially preferred moieties are monocyclic five- or six-membered aromatic hydrocarbons or aromatic heterocycles.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

General synthesis procedures of the polymeric prodrugs

Synthesis of representative examples of polymeric prodrugs according to the present invention is described in the Examples section.

However, prodrugs of the present invention can be prepared in various different fashions. Figure 10 shows general routes for the synthesis of the polymeric prodrugs of the present invention according to formula Ia.

10 In a first method, intermediate (III) is provided by acylating Y_2 of starting material (II) with

$$\text{Nu}-W-\text{Y}_2-\text{C}(=\text{O})-\text{Y}_1$$

For this, X or Nu may have to be protected with a reversible protecting group PG₁. Suitable protecting groups are described in TW Greene, P.G.M. Wuts, Protective groups in organic synthesis, 1999, John Wiley & Sons, 3rd ed.. From intermediate (III) 15 two alternative routes can be used to yield (Iaa). In a first route intermediate (III) is activated by an activating agent such as 4-nitrophenyl chloroformate or disuccinyl carbonate to yield (IV). The amine containing drug molecule is attached to (IV) to yield (V) by displacing the leaving group of the activated intermediate (IV). After 20 deprotection of X such as by treating intermediate (V) with reagents like trifluoroacetic acid or DTT (where applicable) deprotected intermediate (V) is then reacted with polymer R1 to yield the polymeric prodrug (Iaa).

In a second route the polymer R1 is first attached to the intermediate (III) after deprotection of X (where applicable) to form intermediate (VI). After an activation 25 step intermediate (VII) is formed. (VII) is reacted with the amine containing drug molecule to form the polymeric prodrug (Iaa).

In a second method, intermediate (VIII) is provided by activating starting material (II) by an activating agent such as 4-nitrophenyl chloroformate. For this, Y_2 and/or X may 30 have to be protected with a protecting group PG₂ and/or PG₁. Amine containing drug is reacted with intermediate (VIII) to form (IX). In a first route, Y_2 of (IX) is selectively deprotected and acylated to form intermediate (V) which is further processed to (Iaa) as described above. In a second route X is selectively deprotected

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH


5 July 2004

and reacted with polymer R1 to form intermediate (X). Y₂ of (X) is then deprotected and acylated to form the polymeric prodrug (Iaa).

In a third method starting material (II) is reacted with polymer R1 to form intermediate (XI). In one route, intermediate (XI) can be acylated to form intermediate (VI) which processed as described above to form polymeric prodrug (Iaa). In a second route, Y₂ is protected by the protecting group PG₂, activated and reacted with the amine containing drug molecule to form (X). Intermediate (X) is then processed as described above to form the polymeric prodrug (Iaa).

For all methods described, further functional groups such as Y₃ or nucleophiles present in Nu-W may have to be protected with suitable protecting groups.

Polymeric prodrugs according to formula Ib can be prepared by methods described above for prodrugs according to formula Ia using starting material IIb instead of II in figure 10.



It is understood, that linker structures according to the outlined invention and carrying protecting groups or leaving groups as described and used in the synthesis of corresponding polymeric prodrugs are considered within the range of the invention.

Application of the polymeric prodrugs in molecular therapy

One advantage of the described prodrug systems is to enable depot formulations without the need for encapsulation. Until now, many biocompatible materials like hydrogels with large pore sizes could not be used for depot formulations due to their lack of encapsulation properties. From such well-hydrated and mechanically soft biocompatible materials, drugs would be released too fast for most therapeutic applications. In combination with the prodrug linkers described in this invention, the

Ref: CBG00230/GB/F1
Applicant: Complex Biosystems GmbH

5 July 2004

carrier material may be optimized for its biocompatibility properties as the release is solely governed by the linker cleavage kinetics and does not require chemical or enzymatic degradation of the polymer carrier itself.

- 5 A second advantage of the polymeric prodrugs of the present invention is their predominantly non-enzymatic cleavage: the half-life of the prodrug in suitably buffered human blood plasma of pH 7.4 (with aqueous buffer concentration < 50 %) is at least 50 % of the half-life of the prodrug in enzyme-free buffer pH 7.4.
- 10 This feature allows for better predictability and control of release rates after administration to a living organism and reduces interpatient variability.

Release rates are governed by a substantially non-enzymatic chemical reaction which is in turn dependent on the molecular structure of the linker. Systematic or random
15 modifications of the chemical structure, for instance by changing substituents in one or more positions, for instance a masking group in a cascade prodrug, allows for the generation of linkers with differing release rates. It is therefore possible to create a variety of linkers and select those fast or slow cleaving linkers according to the demands posed by a given medicinal or therapeutic application.

20 One embodiment of the invention relates to such cascade-prodrugs where removal of a masking group from an activating moiety triggers a release sequence.

Another advantageous feature which is part of this invention is the attachment of the
25 polymer carrier through a stable covalent bond to an activating moiety involved in a double or cascade prodrug release mechanism. Typically in polymer pro-prodrugs, a bond between the carrier group and the activating moiety is cleaved, this being the release-controlling step, followed by a more rapid molecular rearrangement of the activating moiety and cleavage of a second labile linkage which is located between
30 the activating moiety and the drug. After completion of this sequence, the carrier as well as the activating moiety are released in addition to the drug. Due to the rearrangement, the activating moiety is released in a highly reactive form and may cause direct damage to surrounding biomolecules, or potentially toxic derivatives of this moiety may be formed in vivo. As part of this invention, the activating moiety

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

remains attached to the carrier after drug release and therefore cannot diffuse into the environment. Carrier attachment greatly reduces the reactivity of the moiety and the probability of unwanted side effects.

- 5 In contrast to the enzymatic dependency of masking group removal as described by Antczak, C., Bauvois, B., Monneret, C., Florent J-C, Bioorg Med Chem 9, 2001, 2843), a higher level of control over release rates can be achieved if the masking group has enzyme-independent, self-eliminating properties. The masking groups according to the present invention contain at least one nucleophile Nu. This
- 10 nucleophile, which can be for example a primary, secondary or tertiary amino group can assist the cleavage of the masking group from the activating moiety by intramolecular catalysis or cyclization. For instance the masking group may contain an amine group attached to a four carbon atom chain, which in turn is linked through an ester group to the activating moiety. The free amine may react with this ester in an
- 15 intramolecular fashion, resulting in ester cleavage, amidation and cyclization of the masking group, forming a six-membered ring. Structural features of this masking group such as nucleophilicity of the amine group and ring-forming capacity may be systematically optimized in order to precisely adjust the rate of prodrug cleavage.

- Such intramolecular reactions resulting in unmasking and subsequent rearrangement
- 20 are highly independent from enzymes due to the fact that intramolecular reactions are generally preferred over intermolecular reactions and are shown diagrammatically in Fig.6.

- 25 In another embodiment of the invention, independency of prodrug cleavage from enzyme levels is achieved by providing a prodrug containing a sterically demanding carrier group as is shown in Fig. 7.

- Such encapsulation or sterical protection by the sterically demanding carrier group
- 30 may be conferred by a branched, hyperbranched, crosslinked or self-assembled structure of the carrier polymer. Such polymers tend to form a densely packed molecular volume, as exemplified for instance in dendrimers, dense star polymers or bead-shaped nano- and microparticles or amorphous gels. If the linkage of such a carrier to the drug is located in the interior of such a carrier polymer, the linked drug

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

will be efficiently encapsulated and protected from enzymatic attack. In this case, sterical hindrance by the polymer prevents enzymes from accessing and cleaving the labile linkage.

- 5 In yet another embodiment, enzyme-independent prodrug cleavage is achieved by combining an intramolecular self-eliminating masking group with an encapsulating hyperbranched or crosslinked or self-assembled carrier.

A further advantage of the present invention is the release of an unmodified
10 biologically active moiety. In cases where the biologically active moiety contains further reactive functional groups like amino groups of lysine residues in proteins, an unwanted side reaction between the masking group and the biologically active moiety can occur. The reactive functional groups of the biologically active moiety may react with the masking group, forming a stable covalent bond and resulting in the release of
15 a modified moiety. This potential side reaction is shown schematically in figure 9. The occurrence of such side reactions is shown in the experimental section using polymeric prodrugs which are not part of the present invention with simple masking groups like a pentanoyl residue without a nucleophile Nu present in the masking group. This side reaction is suppressed using polymeric prodrugs according to the
20 present invention with intramolecularly activated masking groups that contain nucleophiles Nu (see experimental section).

Description of the Figures

25

Fig. 1 shows a carrier-linked prodrug.

Fig. 2 shows a cascade prodrug.

Fig. 3 shows cleavage of a prodrug with an amino group.

Fig. 4 shows prodrug cleavage from a carrier which is identical with a masking group.

- 30 Fig. 5 shows cleavage of a cascade prodrug system.

Fig. 6 shows cleavage and intramolecular reaction of the masking group.

Fig. 7 shows cleavage of a prodrug having a sterically demanding carrier group.

Fig. 8 shows cleavage of the masker group by intramolecular cyclisation.

Fig. 9 shows a possible side reaction of polymeric prodrug activation

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Fig.10 shows general synthesis methods

Fig.11 shows mass spectra of prodrug released insulin molecules

Ref: CEG00230/GB/F1
Applicant: Complex Biosystems GmbH

5 July 2004

Examples

Materials

Fmoc-amino acids, resins and PyBOP were purchased from Novabiochem and are
5 named according to the catalogue. Fmoc-Ado-OH was obtained from Neosystem. All
additional chemicals were purchased from Sigma Aldrich. Recombinant human
insulin was from ICN Biomedicals (USA). Maleimide-PEG5k was obtained from
Nektar (USA). 5-(and-6)-carboxyfluorescein succinimidyl ester (mixed isomers) was
obtained from Molecular Probes.

10

Solid phase synthesis reaction medium

Solid phase synthesis was performed on NovaSyn TG Sieber amide resin with a
loading of 0.17 mmol/g or 2-chlorotriyl chloride resin with a loading of 1.4 mmol/g.
Syringes equipped with polypropylene frits were used as reaction vessels.

15

Standard coupling cycle for fmoc-protected amino acids

For fmoc protecting-group removal, the resin was repeatedly (three times, 4 min each)
agitated with 2/2/96 (v/v/v) piperidine/DBU/DMF and repeatedly (six times) washed
with DMF.

20

Coupling of fmoc-protected amino acids to free amino groups on resin was achieved
by agitating the resin with 3 equivalents (eq) of fmoc-amino acid, 3 eq PyBOP and 6
eq DIEA in relation to free amino groups in DMF for 60 min.
Finally, the resin was repeatedly (five times) washed with DMF.

25

Standard cleavage protocol for TentaGel Sieber amide resin

Upon completed synthesis, the resin was washed with DCM, dried in vacuo and
treated repeatedly (five times) with 97/2/1 (v/v) DCM/TES/TFA. After evaporation,
compounds were purified by preparative RP-HPLC (Waters 600).

30

Standard cleavage protocol for 2-chlorotriyl chloride resin

Upon completed synthesis, the resin was washed with DCM, dried in vacuo and
treated two times for 30 minutes with 65/35 (v/v) HFIP/DCM. After combining the
eluates the volatile components were evaporated.

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


Analysis

Mass spectrometry (MS) was performed on a Waters ZQ 4000 ESI instrument and spectra were, if necessary, interpreted by Waters software MaxEnt.

- 5 Size exclusion chromatography was performed using an Amersham Bioscience AEKTA basic system equipped with a Superdex 200 column (Amersham Bioscience).
NMR spectra were recorded on a Bruker AC300.

Overview - synthesis of polymeric prodrugs according to formula Ia with ester-linked masking groups


10



15


20

25



1. NH₂-R₃
2. Mmt deprotection

30




Ref: CBG00230/GB/F1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of polymeric prodrugs according to formula Ia with carbamate-linked
masking groups


a)



Ref: CBG00230/QB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


b)



Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


Synthesis of a polymeric prodrug according to formula Ib with a carbamate-linked masking group



Ref: CBG00230/GB/F1
Applicant: Complex Biosystems GmbH

5 July 2004


Synthesis of compound 2



2

Mmt-chloride (1 eq) and mercaptopropionic acid (1.1 eq) were dissolved in TFA and incubated for 30 min. The solvent was removed under reduced pressure. The product was dissolved in pyridine, diluted in water, acidified by acetic acid and extracted with ether. The ether phase was separated and dried over Na₂SO₄. Solvent was removed under reduced pressure and product 2 was purified by RP-HPLC.

Synthesis of compounds 3a and 3b



3a R1 = H

3b R1 = OM_e

Octopamine hydrochloride 1a or normetanephrine hydrochloride 1b (2 eq), DIEA (4 eq), and PyBOP (1 eq) were dissolved in DMF, 2 (1eq) was added and the mixture was reacted for 50 min at room temperature. After addition of acetic acid (7 eq) products 3a or 3b were purified by RP-HPLC.


3a: MS [M+Na]⁺ = 536 (MW+Na calculated = 536.2 g/mol)

3b: MS [M+Na]⁺ = 566 (MW+Na calculated = 566.2 g/mol)

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of mercaptothiazolide 4




2-Mercaptothiazoline and triethylamine (1.5 eq) were dissolved in dry THF and pentanoyl chloride (1 eq) was added. The mixture was stirred for 1 h at 50 °C under inert gas atmosphere and was allowed to cool to room temperature. 0.5 N aqueous HCl was added and the separated organic phases were dried over Na₂SO₄. After concentration *in vacuo* the residue was purified by silica gel column chromatography using heptane/ethyl acetate (1/1) as mobile phase. Mercaptothiazolide 4 was collected as a viscous yellow oil.

15

4 R_f (heptane/ethyl acetate 1:1) = 0.7

Synthesis of compounds 5a and 5b



25

General synthesis protocol:

1 g 2-chlorotrityl chloride resin (loading 1.6 mmol/g) was incubated for 1 h with 850 mg (2.4 mmol) Fmoc-Ile-OH and 840 µl (4.8 mmol) DIEA in 5 ml 1/1 DCM/DMF. After fmoc removal and washing of the resin with DMF, boc-aminobutyric acid or boc-aminohexanoic acid, respectively, were coupled to 0.5 g resin each according to the standard coupling method. Compounds 5a or 5b, respectively, were cleaved from the resin with 97/1/2 (v/v) DCM/TFA/TES for 45 min. After neutralisation with pyridine, solvents were removed under reduced pressure and 5a or 5b were purified by RP-HPLC.


Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

5a MS $[M+Na]^+ = 339.2$ (MW+Na calculated = 339.4 g/mol)

5b MS $[M+Na]^+ = 367.4$ (MW+Na calculated = 367.5 g/mol)




5 Synthesis of compound 6a



Mercaptothiazolidine 4 (1 eq), phenol 3a (4 eq) and DMAP (4 eq) were refluxed in DCM for 2 h under nitrogen atmosphere. After neutralization with acetic acid, the solvent was removed *in vacuo* and product 6a was purified by RP-HPLC.

6a MS $[M+Na]^+ = 620$ (MW+Na calculated = 620.3 g/mol)

Synthesis of compounds 6b to 6e



6d R1 = OMe
6e R1 = H
32

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

General synthesis protocol:

Carboxylic acid **5a**, **5b** or Z-Lys(Boc)-OH (1 eq), phenol **3b** (1 eq), DIC (1 eq), and DMAP (2 eq) in DMF were reacted for 1h at room temperature. After addition of acetic acid (4 eq) the resulting carboxylic esters **6c**, **6d** or **6b**, respectively, were purified by RP-HPLC.

6e was synthesized as described above using **5b** and **3a** as starting materials.


6b MS $[M+Na]^+ = 928$ (MW+Na calculated = 928.6 g/mol)

6c MS $[M+Na]^+ = 864$ (MW+Na calculated = 864.5 g/mol)

6d MS $[M+Na]^+ = 892$ (MW+Na calculated = 892.6 g/mol)

6e MS $[M+Na]^+ = 862$ (MW+Na calculated = 862.6 g/mol)

Synthesis of compounds **7a** to **7e**



General synthesis protocol:

Alcohols **6a**, **6b**, **6c**, **6d**, or **6e** (1 eq), 4-nitrophenyl chloroformate (10 eq), and DIEA (10 eq) were stirred in dry dioxane for 3 h at room temperature under nitrogen atmosphere. After addition of acetic acid (25 eq) the mixtures were diluted with 7/3

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


Regioselectivity of insulin modification was verified by DTT reduction and MS analysis.

8a MS $[M+2H]^{2+} = 3078.9$; $[M+3H]^{3+} = 2053.2$ $[M+4H]^{4+} = 1540.6$ (MW calculated = 6158 g/mol)

8b MS $[M+2H]^{2+} = 3152.9$; $[M+3H]^{3+} = 2100.6$ $[M+4H]^{4+} = 1575.8$ (MW calculated = 6302 g/mol)

8c MS: $[M+3H]^{3+} = 2110.7$; $[M+4H]^{4+} = 1583.7$; $[M+5H]^{5+} = 1266.6$ (MW calculated = 6330 g/mol)

Synthesis of compounds **8d** to **8g** (N^{εB29}-Fluorescein-N^{αA1}-linker-insulin)



Ref: CBG00230/CB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of N^{εB29}-fluorescein insulin:

80 mg (13.8 μmol) rh-insulin were dissolved in 4 ml 1/1 (v/v) DMF/DMSO and 40 μl DIEA were added. 8 mg (17 μmol) 5-(and-6)-carboxyfluorescein succinimidyl ester were added and the solution was stirred for 30 min at room temperature. 4 ml 5/5/1 (v/v/v) acetonitrile/water/acetic acid were added, product N^{εB29}-fluorescein insulin was purified by RP-HPLC and lyophilized. The conjugation site was verified by reduction of N^{εB29}-fluorescein insulin with 1,4-dithiothreitol, protease digestion and MS analysis.

10 MS: $[M+2H]^{2+} = 3084.0$; $[M+3H]^{3+} = 2054.6$ (MW calculated = 6166 g/mol)

General synthesis protocol:

N^{εB29}-fluorescein insulin in 1/1 (v/v) DMF/DMSO was mixed with a solution of 0.9 eq carbonate 7b, 7c, 7d, or 7e in DMSO. The resulting solutions were adjusted to basic pH with DIEA and stirred for 3 h at RT. RP-HPLC purification gave Mmt-protected intermediates.

After lyophilization, the intermediates were dissolved in 95/5 (v/v) TFA/triethylsilane and stirred for 5 min. Volatiles were removed under nitrogen flow and 8d, 8e, 8f, or 8g were purified by RP-HPLC and lyophilized.

8d MS: $[M+2H]^{2+} = 3364.1$; $[M+3H]^{3+} = 2242.7$; $[M+4H]^{4+} = 1681.5$ (MW calculated = 6724 g/mol)

8e MS: $[M+3H]^{3+} = 2219.2$ $[M+4H]^{4+} = 1665.9$; $[M+5H]^{5+} = 1332.8$ (MW calculated = 6660 g/mol)

8f MS: $[M+3H]^{3+} = 2229.7$ $[M+4H]^{4+} = 1673.3$; $[M+5H]^{5+} = 1337.7$ (MW calculated = 6689 g/mol)

8g MS: $[M+3H]^{3+} = 2218.7$ $[M+4H]^{4+} = 1664.9$ (MW calculated = 6659 g/mol)

30 Synthesis of compounds 9a to 9g (mono-pegylated insulin compounds)

General synthesis protocol:

70 μl of a solution of 8a, 8b, 8c, 8d, 8e, 8f, or 8g (500 μM) in 1/4 (v/v) acetonitrile/water were mixed with 7 μl of a solution of 10 mM maleimide-PEG5k in

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


1/4 (v/v) acetonitrile/water and 10 µl of 0.5 M sodium phosphate buffer (pH 7.0) and incubated for 15 min. Compounds 9a, 9b, 9c, 9d, 9e, 9f, or 9g, respectively, were purified by SEC (column: Superdex 200, flow rate: 0.75 mL/min) using 10 mM HEPES buffer (pH 7.4), 150 mM NaCl, 3 mM EDTA, and 0.005 % Tween as mobile
5 phase. The collected eluates (approximately 1.5 mL) were directly used as such for release rate determination.

9a through 9g: SEC retention time: 19.5 min

10

Synthesis of compounds 11a and 11b

15



11a R₁ = R₂ = Me

11b R₁ = Me, R₂ = H

20 General synthesis protocol:

3,5-Dimethyl-4-hydroxy-acetophenone (5.0 mmol) 10a or 4-hydroxy-3-methyl-acetophenone 10b (0.75 g, 5.0 mmol) and CuBr₂ (1.7 g, 7.5 mmol) were dissolved in 10 ml ethyl acetate and refluxed for 2 h. Solid byproducts were removed by filtration. The filtrate was evaporated and crude products 11a or 11b were purified by RP-
25 HPLC.

11a: Yield 754 mg (62 %)

MS [M+H]⁺ = 243.1/245.1 (MW+H calculated = 244.1 g/mol)


11b: Yield 533 mg (47 %)

30 MS [M+H]⁺ = 229.2/231.1 (MW+H calculated = 230.1 g/mol)

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compounds 12a and 12b



12a R1 = R2 = Me

12b R1 = Me, R2 = H

Synthesis protocol:

- 10 500 mg 11a (2.06 mmol) or 472 mg (2.06 mmol) 11b and 576 mg (4.11 mmol) hexamethylenetetramine were dissolved in 20 ml of trichloromethane and refluxed for 30 min. The solvent was removed in vacuo. 4 ml ethanol and 2 ml of concentrated HCl were added and the slurry was heated to 50 °C for 4 h. The mixture was concentrated in vacuo, diluted with acetonitrile/water and 12a or 12b were purified by
- 15 RP-HPLC.


12a: Yield 547 mg (81 %) as TFA-salt

MS $[M+Na]^+ = 202.2$ (MW+Na calculated = 202.2 g/mol)

12b: Yield 455 mg (70 %) as TFA-salt

20 MS $[M+Na]^+ = 188.2$ (MW+Na calculated = 188.2 g/mol)

Synthesis of compound 13



13

- 500 mg (1.71 mmol) 12a (TFA salt) were dissolved in 10 ml 1/1 (v/v) methanol/water, 129 mg (3.41 mmol) NaBH₄ were added and the mixture was stirred
- 30 for 30 min at RT. 0.5 ml of acetic acid were added and 13 was purified by RP-HPLC.

13: Yield 313 mg (62 %) as TFA-salt


MS $[M+Na]^+ = 204.2$ (MW+Na calculated = 204.2 g/mol)

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

NMR (300 MHz, DMSO- d_6) δ [ppm] = 8.25 (s, 1H, Phenol), 7.84 (bs, 3H, NH_3^+), 6.89 (s, 2H, CH_{ar}), 5.85 (d, 1H, Hydroxyl, $J=3.7$ Hz), 4.62 (m, 1H, CH_{Benzyl}), 2.93 (m, 1H, CH_a), 2.80 (m, 1H, CH_b), 2.17 (s, 6H, CH_3).

5 Synthesis of compounds 14a to 14c



- 10 **14a** R1 = R2 = Me, R7 = H
14b R1 = R2 = H, R7 = Me
14c R1 = OMe, R2 = H, R7 = Me

13 (TFA salt, 159 mg, 0.541 mmol), synephrine (335 mg, 2.00 mmol), or
 15 metanephrine (HCl salt, 281 mg, 1.20 mmol) were coupled to compound 2 as described for compound 3a to yield 14a, 14b, or 14c, respectively.

14a: Yield 254 mg (87 %)

MS $[M+Na]^+ = 564.7$ (MW+Na calculated = 564.3 g/mol)


20 **14b:** Yield 760 mg (72 %)

MS $[M+Na]^+ = 550.2$ (MW+Na calculated = 550.3 g/mol)

14c: Yield 530 mg (80 %)

MS $[M+Na]^+ = 580.4$ (MW+Na calculated = 580.4 g/mol)

25 Synthesis of compounds 15c, 15d and 15f



30

General synthesis protocol:

1 g 2-chlorotriptyl chloride resin (loading 1.4 mmol/g) was incubated for 1 h with N,N' -dimethylpropane-1,3-diamine (for synthesis of 15c), or N,N' -diethylpropane-

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


1,3-diamine (for synthesis of 15d) or N,N'-dimethyl-ethane-1,2-diamine (4 eq) (for synthesis of 15f) in DCM. After washing of the resin with DMF, amines were acetylated with 1/1/2 (v/v/v) acetic anhydride/pyridine/DMF for 14 h. The resin was washed with THF and dried. LiAlH₄ (1 M in THF, 4 eq) was added dropwise to the suspended resin in THF. The resulting suspension was stirred for 3 h at 45°C under nitrogen atmosphere. After cooling, aqueous Rochelle's salt solution was added and the resin was separated and dried. Compounds were cleaved from resin with 2/1 (v/v) HFIP/DCM (2 x 30 min). The volatile components were evaporated and the products 15c, 15d or 15f were used in the following steps without further purification.

15c MS [M+H]⁺ = 131.2 (MW = 130.1 g/mol)

15d MS [M+H]⁺ = 159.2 (MW = 158.1 g/mol)

15f MS [M+H]⁺ = 117.1 (MW = 116 g/mol)

Synthesis of compounds 16a to 16f



16a R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl, R7 = H

16b R1 = OMe, R2 = H, R5 = Et, R6 = 2-(diethylamino)ethyl, R7 = H

16c R1 = OMe, R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

16d R1 = R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

16e R1 = OMe, R2 = H, R5 = Et, R6 = 3-(diethylamino)propyl, R7 = Me

16f R1 = R2 = H, R5 = Et, R6 = 3-(diethylamino)propyl, R7 = Me

14a (120 mg, 0.222 mmol) was dissolved in 1.5 ml of dry THF. p-Nitrophenyl-chloroformate (45 mg, 0.222 mmol) and DIEA (113 µl, 0.665 mmol) were added and the mixture was stirred for 30 min at RT. 15a (N,N,N'-trimethyl-ethylene-1,2-diamine) (72 µl, 0.554 mmol) was added and stirring was continued for 30 min. The solvent was removed in vacuo, 100 µl of AcOH were added and 16a was purified by RP-HPLC.

Ref: CEG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

16b was synthesized as described above from 3b (80 mg, 0.15 mmol) and 15b (N,N,N'-triethyl-ethylene-1,2-diamine) (55 mg, 0.38 mmol).

16c or 16d were synthesized as describe above from 14c (56 mg, 0.1 mmol) or 14b (53 mg, 0.1 mmol), respectively, and diamine 15c.

- 5 16e or 16f were synthesized as described above from 14c (56 mg, 0.1 mmol) or 14b, respectively, (53 mg, 0.1 mmol) and diamine 15d.

16a: Yield 120 mg (69 %) as TFA salt

MS $[M+Na]^+ = 692.4$ (MW+Na calculated = 692.9 g/mol)

- 10 16b: Yield 48 mg (40 %) as TFA salt

MS $[M+Na]^+ = 736.3$ (MW+Na calculated = 736.4 g/mol)

16c: Yield 8 mg (10 %) as TFA salt

MS $[M+Na]^+ = 736.4$ (MW+Na calculated = 736.4 g/mol)

16d: Yield 20 mg (25 %) as TFA salt

- 15 MS $[M+Na]^+ = 706.3$ (MW+Na calculated = 706.3 g/mol)

16e: Yield 2 mg (3 %) as TFA salt


MS $[M+Na]^+ = 764.6$ (MW+Na calculated = 764.4 g/mol)

16f: Yield 6 mg (8 %) as TFA salt

MS $[M+Na]^+ = 734.4$ (MW+Na calculated = 734.3 g/mol)

20

Synthesis of compound 17



25

17

Amine 12b (TFA salt) was coupled to compound 2 as described for compound 3a.


- 30 17: Yield 608 mg (74 %)

MS $[M+Na]^+ = 548.3$ (MW+Na calculated = 548.7 g/mol)

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compounds 18a and 18b



18a R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl

18b R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl

10 383 mg (0.729 mmol) of 17 were reacted with p-nitrophenyl-chloroformate and N,N,N'-trimethyl-propane-1,3-diamine (15e) or 15f, respectively, to yield 18a or 18b as described for compound 16a.


18a: Yield 287 mg (50 %) as TFA salt

MS $[M+Na]^+ = 690.7$ (MW+Na calculated = 690,9 g/mol)

15 **18b:** Yield 148 mg (26 %) as TFA salt

MS $[M+Na]^+ = 690.9$ (MW+Na calculated = 690,9 g/mol)

Synthesis of compounds 16g and 16h



16g R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl

16h R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl

25 **18a** (287 mg, 0.367 mmol, TFA salt) was dissolved in 5 ml methanol, NaBH₄ (41 mg, 1.07 mmol) was added and the mixture was stirred for 30 min at RT. 0.5 ml of acetic acid were added and 16g was purified by RP-HPLC.

18b (8 mg, 0.010 mmol, TFA salt) was reacted as described above to yield 16h.

30 **16g:** Yield 201 mg (70 %) as TFA-salt

MS $[M+Na]^+ = 692.7$ (MW+Na calculated = 692.9 g/mol)


16h: Yield 6 mg (77 %) as TFA-salt

MS $[M+Na]^+ = 692.7$ (MW+Na calculated = 692.9 g/mol)

Ref: CRG00230/GB/F1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compounds 19a to 19h



19a R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl, R7 = H

19b R1 = OMe, R2 = H, R5 = Et, R6 = 2-(diethylamino)ethyl, R7 = H

19c R1 = OMe, R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

10 19d R1 = R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me

19e R1 = OMe, R2 = H, R5 = Et, R6 = 3-(diethylamino)propyl, R7 = Me

19f R1 = R2 = H, R5 = Et, R6 = 3-(diethylamino)propyl, R7 = Me

19g R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl, R7 = H

19h R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl, R7 = H

15

Carbonates 19a to 19h were synthesized from 16a to 16h, respectively, as described for compound 7a.

19a: Yield 98 mg (72 %) as TFA-salt

20 MS $[M+Na]^+$ = 857.8 (MW+Na calculated = 858.0 g/mol)

19b: Yield 6 mg (11 %) as TFA-salt

MS $[M+Na]^+$ = 901.8 (MW+Na calculated = 901.5 g/mol)

19c: Yield 1 mg (15 %) as TFA-salt

MS $[M+Na]^+$ = 901.4 (MW+Na calculated = 901.5 g/mol)

25 19d: Yield 8 mg (29 %) as TFA-salt

MS $[M+Na]^+$ = 871.4 (MW+Na calculated = 871.4 g/mol)

19e: Yield 0.3 mg (18 %) as TFA-salt

MS $[M+Na]^+$ = 929.4 (MW+Na calculated = 929.5 g/mol)

19f: Yield 4 mg (45 %) as TFA-salt

30 MS $[M+Na]^+$ = 899.7 (MW+Na calculated = 899.6 g/mol)

19g: Yield 6 mg (6 %) as TFA-salt

MS $[M+Na]^+$ = 857.8 (MW+Na calculated = 858.0 g/mol)


19h: Yield 0.8 mg (11 %) as TFA-salt

MS $[M+Na]^+$ = 857.7 (MW+Na calculated = 858.0 g/mol)

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compounds 20a to 20f




- 10 20a R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl, R7 = H
 20b R1 = OMe, R2 = H, R5 = Et, R6 = 2-(diethylamino)ethyl, R7 = H
 20c R1 = OMe, R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me
 20d R1 = R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me
 20e R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl, R7 = H
 15 20f R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl, R7 = H

Insulin derivatives 20a, 20b, 20c, 20d, 20e, or 20f were synthesized from 19a, 19b, 19c, 19d, 19g, or 19h respectively, as described for compound 8a.

- 20 20a MS $[M+3H]^{3+} = 2077.3$ $[M+4H]^{4+} = 1559.2$ (MW calculated = 6231.3 g/mol)
 20b MS $[M+3H]^{3+} = 2093.0$ $[M+4H]^{4+} = 1569.6$ (MW calculated = 6274 g/mol)
 20c MS $[M+3H]^{3+} = 2090.8$ $[M+4H]^{4+} = 1568.7$ (MW calculated = 6274 g/mol)
 20d MS $[M+3H]^{3+} = 2081.3$ $[M+4H]^{4+} = 1561.8$ (MW calculated = 6244 g/mol)
 20e MS $[M+3H]^{3+} = 2077.1$ $[M+4H]^{4+} = 1558.2$ (MW calculated = 6231.3 g/mol)
 25 20f MS $[M+3H]^{3+} = 2076.7$ $[M+4H]^{4+} = 1559.3$ (MW calculated = 6231.3 g/mol)

Synthesis of compounds 21a to 21f (mono-pegylated insulin derivatives)



Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


- 21a R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl, R7 = H
 21b R1 = OMe, R2 = H, R5 = Et, R6 = 2-(diethylamino)ethyl, R7 = H
 21c R1 = OMe, R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me
 5 21d R1 = R2 = H, R5 = Me, R6 = 3-(N-ethyl-N-methylamino)propyl, R7 = Me
 21e R1 = R5 = Me, R2 = H, R6 = 3-(dimethylamino)propyl, R7 = H
 21f R1 = R5 = Me, R2 = H, R6 = 2-(N-ethyl-N-methylamino)ethyl, R7 = H

- Insulin derivatives 21a, 21b, 21c, 21d, 21e, or 21f were synthesized from compound
 10 20a, 20b, 20c, 20d, 20e, or 20f, respectively, as described for compound 9a.

21a through 21f: SBC retention time: 19.5 min

Synthesis of compounds 23a and 23b

15



- 23a R1 = Me, R2 = H
 20 23b R1 = R2 = Me

- o-Cresol (22a) or 2,6-dimethylphenol (22b) (1 eq), succinic anhydride (1 eq), and
 AlCl₃ (3 eq) in nitrobenzene were heated to 100 °C for 1 h. The reaction mixture was
 poured on HCl/ice and extracted with ether. The organic layer was extracted with 1 N
 25 NaOH and the aqueous layer was acidified with concentrated HCl. The aqueous layer
 was extracted with ether and the ether was evaporated. 23a or 23b were purified by
 RP-HPLC.

- 23a: Yield 552 mg (31 %)
 30 MS [M+Na]⁺ = 231.0 (MW+Na calculated = 231.2 g/mol)
 NMR (300 MHz, DMSO-d₆) δ[ppm] = 12.05 (bs, 1H, CO₂H), 10.23 (s, 1 H, phenol
 OH), 7.74 (s, 1H, CH_{ar}), 7.7 (d, 1H, CH_{ar}, ³J_{H,H} = 8.4 Hz), 6.86 (d, 1H, CH_{ar}, ³J_{H,H} =

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

8.4 Hz), 3.13 (t, 2H, C(O)CH₂, ³J_{H,H} = 6.4 Hz), 2.53 (t, 2H, CH₂CO₂, ³J_{H,H} = 6.4 Hz),
2.16 (s, 3H, CH₃)

23b: Yield 166 mg (15 %)

MS [M+Na]⁺ = 245.4 (MW+Na calculated = 245.2 g/mol)

5

Synthesis of compound 24




1.85 g (16.02 mmol) cysteamine hydrochloride were dissolved in 15 ml of TFA and
2.47 g (8.01 mmol) MmtCl were added. After stirring the mixture at RT for 20 min
the solvent was evaporated in vacuo. The residue was dissolved in diethyl ether and
extracted with saturated NaHCO₃, 1N H₂SO₄ and brine. The solvent was evaporated
and **24** was purified by RP-HPLC.

24: Yield 1.11 g (30 %) as TFA salt

TLC (AcOEt/Et₃N 99/1), R_f = 0.24

Synthesis of compounds 25a and 25b



25a R1 = Me, R2 = H

25b R1 = R2 = Me

25

23a or **23b** (1eq), HOBT (1.1 eq) and DIC (1 eq) were dissolved in DMF and stirred
at RT for 30 min. **24** (TFA salt 1 eq) and DIEA (3 eq) were added and the solution
was stirred for 60 min. Acetic acid was added and **25a** or **25b** was purified by RP-
HPLC.

30

25a: Yield 552 mg (25 %)

MS [M+Na]⁺ = 562.7 (MW+Na calculated = 562.7 g/mol)


25b: Yield 15 mg (40 %)

MS [M+Na]⁺ = 576.6 (MW+Na calculated = 576.6 g/mol)

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compounds 26a and 26b



26a R1 = Me, R2 = H, R5 = R6 = 3-(dimethylamino)propyl

26b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl

10 **26a** 267 mg (0.495 mmol) **25a** was reacted with p-nitrophenylchloroformate and N-(3-dimethylamino-propyl)-N',N'-dimethyl-propane-1,3-diamine (**15g**) to yield **26a** as described for compound **16a**.

26b was synthesized as described above using 15 mg **25b** and N,N,N'-trimethyl-ethane-1,2-diamine (**15a**).

15 **26a**: Yield 282 mg (58 %) as double TFA salt


MS [M+Na]⁺ = 775.2 (MW+Na calculated = 776.0 g/mol)

26b: Yield 17 mg (70 %) as TFA salt

MS [M+Na]⁺ = 704.5 (MW+Na calculated = 704.6 g/mol)

20

Synthesis of compounds 27a and 27b



27a R1 = Me, R2 = H, R5 = R6 = 3-(dimethylamino)propyl

27b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl

30 **26a** (272 mg, 0.277 mmol, double TFA salt) was dissolved in 5 ml methanol, NaBH₄ (42 mg, 1.09 mmol) was added and the mixture was stirred for 30 min at RT. 0.5 ml of acetic acid were added and **27a** was purified by RP-HPLC.

Alcohol **27b** was synthesized likewise from **26b** (17 mg, 25 μmol, TFA salt).

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

27a: Yield 142 mg (52 %) as double TFA salt


MS $[M+Na]^+ = 777.9$ (MW+Na calculated = 778.0 g/mol)

27b: Yield 6 mg (40 %) as TFA salt

5 MS $[M+Na]^+ = 706.5$ (MW+Na calculated = 706.6 g/mol)

Synthesis of compounds 28a and 28b

10



15 **28a** R1 = Me, R2 = H, R5 = R6 = 3-(dimethylamino)propyl

28b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl

20 Carbonates **28a** or **28b** were synthesized from **27a** or **27b**, respectively, as described for compound **7a**.

28a: Yield 1 mg (29 %)


MS $[M+Na]^+ = 942.9$ (MW+Na calculated = 943.2 g/mol)

28b: Yield 1.5 mg (19 %)

25 MS $[M+Na]^+ = 871.6$ (MW+Na calculated = 871.7 g/mol)

Synthesis of compounds 29a and 29b

30



Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

29a R1 = Me, R2 = H, R5 = R6 = 3-(dimethylamino)propyl

29b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl


- 5 Insulin derivatives **29a** or **29b** were synthesized from **28a** or **28b**, respectively, as described for compound **8a**.

29a MS $[M+3H]^{3+} = 2105.8$ $[M+4H]^{4+} = 1580.2$ (MW calculated = 6316.4 g/mol)

29b MS $[M+3H]^{3+} = 2081.8$ $[M+4H]^{4+} = 1562.4$ (MW calculated = 6244 g/mol)

10

Synthesis of mono-pegylated insulin derivatives **30a** and **30b**



30a R1 = Me, R2 = H, R5 = R6 = 3-(dimethylamino)propyl

30b R1 = R2 = R5 = Me, R6 = 2-(dimethylamino)ethyl

- 25 Insulin derivatives **30a** or **30b** were synthesized from **29a** or **29b**, respectively, as described for compound **9a**.


30a and **30b**: SEC retention time: 19.5 min

30


Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

R2 =




R1 =



Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compound 31



31

5

31 was obtained according to the standard solid-phase synthesis protocol. The amino acids Fmoc-Dpr(Boc)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Ado-OH, and Fmoc-Dpr(Fmoc)-OH were coupled to NovaSyn TG Sieber amide resin. After final fmoc removal the resin was agitated with 5 eq maleimidopropionic acid and 5 eq DIC in relation to amino groups in DMF for 30 min. 31 was cleaved from resin with TFA/TES/water 95/3/2 (v/v/v). After evaporation of solvent, product 31 was purified by RP-HPLC.


10

MS: $[M+H]^+ = 2494.6$ (MW calculated = 2495.4 g/mol)

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compound 32



32

Compound **32** was obtained according to the standard solid-phase synthesis protocol. The amino acids Fmoc-Cys(Mmt)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Ado-OH, and Fmoc-Dpr(Fmoc)-OH were coupled to NovaSynTG Sieber amide resin.


After final fmoc removal the resin was agitated with 3 eq Boc-aminoxylacetic acid, 3 eq DIC, and 3 eq HOBT in relation to amino groups in DMF for 30 min. **32** was cleaved from resin with DCM/TFA/TES 97/1/2 (v/v/v). After addition of 0.8 eq pyridine in relation to TFA the solvent was evaporated and product **32** was purified by RP-HPLC.

MS: $[M+H]^+ = 2688.2$ g/mol (MW calculated = 2688.8 g/mol)

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compound 33



33

- 5 Compound 33 was obtained according to the standard solid-phase synthesis protocol. The amino acids Fmoc-Dpr(ivDde)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Dpr(Fmoc)-OH, Fmoc-Lys(Fmoc)-OH, and Fmoc-Ado-OH were coupled to NovaSyn TG Sieber amide resin.

10 After final fmoc removal the resin was agitated with 3 eq 3,6,9-trioxadecanoic acid, 3 eq PyBOP, and 6 eq DIEA in relation to amino groups in DMF for 60 min.

To cleave the ivDde protecting group, the resin was treated three times with 2 % hydrazine in DMF. After washing, 3 eq Fmoc-Ser-OH was coupled with 3eq DIC and 3 eq HOBt for 30 min. After final fmoc removal resin was washed and the product was cleaved from resin with DCM/TFA/TES 88/10/2 (v/v/v). Solvent was evaporated and the residue was oxidized with 10 eq sodium periodate in 3/2 (v/v) 0.1 M sodium phosphate pH 7/acetonitrile for 15 min to yield 33. Product 33 was purified by RP-HPLC and lyophilized.


15

MS: $[M+H]^+ = 3372.1$ g/mol (3372.8 g/mol)

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compound 34



Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

6 mg (2.4 μ mol) of compound 31 were dissolved in 1 ml 2/1 (v/v) acetonitrile/0.1 M sodium phosphate buffer pH 7 and 65 mg (24.2 μ mol) of compound 32 were added. The solution was stirred at room temperature for 2 h and then the product purified by
5 RP-HPLC and lyophilized (yield: 45 mg (78 %)).

The lyophilized product (45 mg) was dissolved in 0.5 ml DMF and 10 μ l DIEA were added. 5 mg (30 μ mol) 3-maleimidopropionic acid and 4.7 μ l (30 μ mol) DIC in 150 μ l DMF were added and the reaction mixture was stirred at room temperature for 20
10 min, the product purified by RP-HPLC and lyophilized.

The lyophilized product was incubated for 10 min in 95/5 (v/v) TFA/water and then the solvent was removed in a stream of nitrogen. Product 34 was purified by RP-HPLC and lyophilized (overall yield for all three steps: 20 mg (47 %)).
15

MS: 17700 – 18200 (broad peak) (MW calculated = 17749 g/mol)

Synthesis of compound 9h

20

1.5 mg (225 nmol) 8g and 5 mg (280 nmol) 34 were mixed, dissolved in 300 μ l 2/1 (v/v) 0.1 M sodium phosphate buffer pH 7/acetonitrile and incubated for 15 min at room temperature. The product was purified by RP-HPLC and lyophilized. (yield 4 mg, 160 nmol, 70 %)

25


The lyophilized product was dissolved in 200 μ l 0.1 M sodium citrate buffer pH 1.5 and 69 mg (20.5 μ mol) 33 in 200 μ l 2/1 (v/v) acetonitrile/sodium citrate buffer pH 1.5 were added. The mixture was stirred at room temperature for 24 h and product 9h was purified by size exclusion chromatography (column: Superdex 200, buffer: 10 mM
30 HEPES pH 7.4, 0.005 % Tween-20, 3mM EDTA, flow rate: 0.75 ml/min)

SEC elution time: 15 min

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compound 38



250 mg (0.35 mmol) 2-chlorotriyl chloride resin (loading 1.4 mmol/g) was incubated
10 for 1.5 h with 308 mg (4 eq., 1.4 mmol) 4,7,10-trioxatridecane-1,13-diamine in 4 ml
DCM to yield 35. The resin was washed with DCM and dried. 107 mg (0.7 mmol)
HOBt, 110 µl (0.7 mmol) DIC, and 150 mg (0.9 mmol) 5-formyl salicylic acid (36) in
3 ml DMF were added and the resulting suspension was stirred for 1 h at RT to yield
37. After washing with DCM and THF, the resin was suspended in 6 mL THF and 3
15 ml (3 mmol) BH₃ THF (1 M in THF, 8.5 eq.) were added dropwise. The reaction
mixture was stirred for 18 h at 45°C under nitrogen atmosphere. After cooling 4 ml
THF, 0.8 ml DIEA and 1.6 ml MeOH were added successively. 210 mg (0.84 mmol)
I₂ (as a concentrated THF solution) were added and the suspension was stirred for 1 h.
The resin was repeatedly washed (three times each) with THF, DMF, MeOH, and
20 DCM. The dried resin was reacted with 107 mg (0.7 mmol) HOBt, 110 µl (0.7 mmol)
DIC, and 55 µL (0.9 mmol) AcOH in 3 ml DMF for 1h. After washing of the resin
with DMF and DCM compound 38 was cleaved from resin with 2/1 (v/v) HFIP/DCM
(2 x 30 min). The volatile components were evaporated and the product 38 was used
in the following step without further purification.

25

38: Yield 29 mg (20 %) as TFA salt


MS [M+Na]⁺ = 421.4 (MW+Na calculated = 421.5 g/mol)

30

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compound 39




24 mg (0.06 mmol) 38, 31 mg (0.06 mmol) PyBOP, 32 μ l (0.18 mmol) DIEA, and 23
10 mg (0.06 mmol) 2 in 0.5 mL DMF were reacted for 50 min at room temperature. After
addition of 50 μ l acetic acid product 39 was purified by RP-HPLC.

39: Yield 7 mg (15 %) as TFA salt

15 MS $[M+Na]^+ = 781.3$ (MW+Na calculated = 781.6 g/mol)

Synthesis of compound 40



39 (7 mg, 9 μ mol) was dissolved in 200 μ l of dry THF. p-Nitrophenylchloroformate
(2.0 mg, 10 μ mol) and DIEA (4.4 μ l, 25 μ mol) were added and the mixture was stirred
for 30 min at RT. N,N,N'-Triethylethylenediamine (15b) (18 μ l, 0.1 mmol) was
30 added and stirring was continued for 30 min. The solvent was removed *in vacuo*, 10
 μ l of AcOH were added and 40 was purified by RP-HPLC.

40: Yield 1 mg (11 %) as TFA salt

MS $[M+Na]^+ = 951.1$ (MW+Na calculated = 951.8 g/mol)


Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of compound 41

5

10



Carbonate **41** was synthesized from **40** as described for compound **7a**

15

41: Yield 0.4 mg as TFA salt

MS $[M+Na]^+ = 1116.8$ (MW+Na calculated = 1116.9 g/mol)

20


25

30

Ref: CB000230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Synthesis of rh-insulin loaded PEGA hydrogel 45



Maleimide derivatization of polyacrylamide based hydrogel (PEGA):

NH₂-PEGA hydrogel beads with 0.4 mmol/g loading and 150-300 μm bead size were purchased from Novabiochem.

30 2.5 g methanol-wet NH₂-PEGA-hydrogel (0.4 mmol/g NH₂-loading) was weighed into a syringe equipped with a polypropylene frit. Maleimide loading was adjusted by acylation employing a mixture of activated maleimidopropionic acid and acetic acid as described in the following. The hydrogel was washed 5 times with DMF and reacted with 13.5 mg (0.08 mmol) 3-maleimidopropionic acid, 115.2 μl (1.92 mmol)

5 July 2004

5 30 mg of maleimide derivatized resin **44** (loading 16 $\mu\text{mol/g}$) was reacted with 3 mg of compound **20b** (480 nmol, 1.06 eq) in 600 μl 20/80 (v/v) acetonitrile/50 mM phosphate buffer (pH 7.4) for 10 min to give rh-insulin loaded hydrogel **45**. The hydrogel **45** was washed 5 times with 50/50 (v/v) acetonitrile/water and three times with acetonitrile and dried under vacuum.

10



20

25

Ref: CEG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

1.5 mg **8c** was dissolved in 25/75 (v/v) acetonitrile/50 mM phosphate buffer pH 7.4 and reacted with 10.8 mg maleimide derivatized hydrogel for 10 min. The rh-insulin loaded hydrogel **46** was washed five times with 50/50 (v/v) acetonitrile/water and three times with acetonitrile and dried under vacuum.

5

Release of insulin or fluorescein-insulin from conjugates in buffer pH 7.4

Release of (fluorescein)-insulin from (fluorescein)-insulin conjugates **9a** to **9h**, **21a** to **21f**, **30a** and **30b** was effected by linker hydrolysis in aqueous buffer pH 7.4.

- 10 Collected SEC eluates of (fluorescein) insulin conjugates (see above) were incubated at 37 °C and samples were taken at time intervals and analyzed by RP-HPLC (insulin conjugates) or SEC (fluorescein insulin conjugates) and UV detection. Peaks correlating with the retention time of native insulin or fluorescein-insulin, respectively, were integrated and plotted against incubation time, and curve-fitting
15 software was applied to estimate the corresponding halftime of release.

Release of insulin from hydrogel conjugates **45** and **46**

- 4 mg of **45** or 2 mg **46** were weighed into a test tube and incubated with 1 ml 10 mM
20 HEPES buffer pH 7.4, 150 mM NaCl, 0.005 % Tween at 37°C. 45 µl samples were taken at different time intervals and quantitatively analyzed for rh-insulin by a RP-HPLC assay. The rh-insulin peak was integrated and rh-insulin concentration was obtained from a standard curve. A first order release kinetic was fitted to the data points to give the linker half life.

25

MS-Analysis of released insulin from compound **9a**, **9b** and **30a**

- Samples of buffer released insulin (see above) were analyzed by mass spectrometry. Figure 11 shows the mass spectra of released insulin from compound **9a**, **9b**, and **30a**.
30 The mass spectrum of insulin released from compound **9a** clearly shows a major side product (indicated by arrows), corresponding to the irreversibly pentanoyl-modified insulin. In this case, removal of the pentanoyl masking group was not by hydrolysis but by acyl transfer to the insulin. The mass spectrum of insulin released from compound **9b** and **30a** shows no modification.

Ref: CBG00230/CB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Release of fluorescein-insulin from conjugate 9d and 9e in 80 % human plasma

5 Release of fluorescein-insulin from 9d or 9e was effected by hydrolysis in 80 % human plasma in 20 mM HEPES pH 7.4 at 37 °C. Samples were taken at time intervals and analyzed by SEC and UV detection. Peaks correlating with the retention time of fluorescein-insulin were integrated and plotted against incubation time, and curve-fitting software was applied to estimate the corresponding halftime of release.

10 Table: Polymeric prodrug hydrolysis

compound	t _{1/2} buffer pH 7.4	t _{1/2} human plasma
9a	40 h	nd
9b	55 h	nd
9c	4.5 d	nd
9d	7 h	4 h
9e	55 h	30 h
9f	90 h	nd
9g	37 h	nd
9h	88 h	nd
21a	64 d	nd
21b	8 d	nd
21c	52 d	nd
21d	29 d	nd
21e	100 d	nd
21f	83 h	nd
30a	17 d	nd
30b	> 70 d	nd
45	7 d	nd
46	4 d	nd

nd = not determined

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Abbreviations:

	Boc	t-butyloxycarbonyl
	DBU	1,3-diazabicyclo[5.4.0]undecene
	DCM	dichloromethane
5	(iv)Dde	1-(4,4-dimethyl-2,6-dioxo-cyclohexyliden)3-methyl-butyl
	DIC	diisopropylcarbodiimide
	DIEA	diisopropylethylamine
	DMAP	dimethylamino-pyridine
	DMF	N,N-dimethylformamide
10	DMSO	dimethylsulfoxide
	Dpr	diaminopropionic acid
	EDTA	ethylenediaminetetraacetic acid
	Et	ethyl
	eq	stoichiometric equivalent
15	fmoc	9-fluorenylmethoxycarbonyl
	Fmoc-Ado-OH	Fmoc-8-amino-3,6-dioxaoctanoic acid
	HFIP	hexafluoroisopropanol
	HEPES	N-(2-hydroxyethyl) piperazine-N'-(2-ethanesulfonic acid)
	HOBt	N-hydroxybenzotriazole
20	LCMS	mass spectrometry-coupled liquid chromatography
	Mal	maleimidopropionyl
	Me	methyl
	Mmt	4-methoxytrityl
	MS	mass spectrum
25	MW	molecular mass
	PyBOP	benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate
	RP-HPLC	reversed-phase high performance liquid chromatography
	RT	room temperature
30	SEC	size exclusion chromatography
	Suc	succinimidopropionyl
	TES	triethylsilane
	TFA	trifluoroacetic acid

Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

THF	tetrahydrofurane
UV	ultraviolet
Z	benzyloxycarbonyl

5



Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Fig. 1




Fig. 2





Fig. 3





Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Fig. 4






Fig. 5



Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004

Fig. 6



\curvearrowright Site of Chemical Transformation
(Non-Enzymatic Cleavage)

Fig. 7





Fig. 8






Ref: CBG00230/GB/P1
Applicant: Complex Biosystems GmbH

5 July 2004


Fig. 9



5 July 2004

Ref: CBG00230/GB/PJ
Applicant: Complex Biosystems GmbH


Fig. 10



Ref: CBG03200GB/P1
Applicant: Compex Biosystems GmbH

5 July 2004

Fig. 11





1